

The Absence of Superfluid Response in the Optical Conductivity of the Edge Region of Optimally Doped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ Crystals

J.J. Tu, C.C. Homes, M. Strongin (BNL), D.N. Basov (UCSD), and G.D. Gu (UNSW, Australia)

Abstract No. Tu5437

Beamline(s): U10A

Introduction: The edge regions of high- T_c superconductors have been probed extensively by tunneling experiments on the ac, bc and (110) faces of these materials into the ab-plane, particularly by Greene and co-workers. One particularly interesting result is the zero bias conductance peak (ZBCP) observed when tunneling on the ac or bc-faces coupled with the absence of a gap feature in the tunneling spectra. This is contrasted with the observation of a weaker ZBCP when tunneling on the (110) face with the appearance of a superconducting gap. When tunneling on the ab-plane into the c-axis, no ZBCP is observed. In this study, we extend the optical conductivity measurements to the same regions near the edges of ab-plane crystals in an effort to examine if the edge regions in the high- T_c d-wave superconductors have any unusual properties that are different from the bulk.

Methods and Materials: Large optimally doped BSCCO single crystals are grown using a traveling-surface-floating-zone (TSFZ) method. Temperature dependent reflectance is measured in a near-normal-incidence arrangement from $\sim 50 \text{ cm}^{-1}$ to over $16,000 \text{ cm}^{-1}$ on a Bruker IFS 66v/S. The absolute reflectivity is determined by an *in situ* evaporation technique and the optical conductivities are then determined from a Kramers-Kronig analysis.

Results: In **Figure 1**, the temperature dependent ab-plane conductivity data is shown for a single crystal BSCCO sample with E-vector parallel to the a-axis. Because of the high signal-to-noise ratio achieved using our *in situ* evaporation technique, two optic phonons at 477 cm^{-1} and 630 cm^{-1} have been observed in σ_1 and σ_2 for the first time in optical conductivity measurements on BSCCO. The superfluid response is observed in the ab-plane conductivities below T_c as σ_1 decreases with temperature according to the Ferrel-Glover-Tinkham sum rule accompanied by a simultaneous increase in σ_2 . In **Figure 2**, we have a set of data to show that with a coarser polishing finish ($1\mu\text{m}$ diamond film) the conductivity of the bc-plane in the normal state is drastically reduced. There is a large spectral feature at 627 cm^{-1} that appears as an anti-resonance dip in σ_1 . In the insert of **Figure 2(a)**, room temperature reflectance data is given with the E-vectors parallel to the b-axis and the c-axis on this bc-face. Strong anisotropy is observed in the reflectance curves, which leads us to conclude that there is no significant contamination by c-axis phonons in the conductivity data with E-vector parallel to b-axis. The temperature dependence of σ_1 and σ_2 is characteristically different from the ab-plane data and in particular there is no evidence of superfluid response in σ_1 or σ_2 . However, the superfluid contribution to optical conductivity is measured in the surface region of the (110)-plane where the E-vector is parallel to an anti-nodal direction.

Conclusions: We have observed characteristically different behavior in the ac and bc-plane optical conductivities of optimally doped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ single crystals as compared to the (110)-plane conductivity below T_c . This apparent asymmetry implies that the surface region in the high- T_c d-wave superconductors has unusual properties that are different from the bulk.

Acknowledgments: We thank T. Valla, Z. Yusof, P.D. Johnson, G.L. Carr, A. Millis, V.J. Emery and T. Timusk for helpful discussions. The work was supported by the U.S. Department of Energy under Contract No. DE-AC02-98CH10886. Research undertaken at NSLS was supported by the U.S. DOE, Division of Materials and Chemical Sciences.




Figure 1. ab-plane optical conductivities of a BSCCO single crystal.




Figure 2. bc-plane optical conductivities of a BSCCO single crystal.